


Your important health information

Pregnancy related carpal tunnel syndrome



What is the carpal tunnel?

The carpal tunnel is formed by the bones of the wrist and is covered by a ligament made of strong connective tissue. This tunnel provides a small passage into the hand for the nerves and tendons travelling to the fingers.

What is carpal tunnel syndrome?

Carpal tunnel syndrome is a painful condition caused by compression of the median nerve within the carpal tunnel. It occurs when the structures in or around the nerve become inflamed and swollen, resulting in a pressure build up in the tunnel itself. Fluid retention in your hands during pregnancy can cause this.

Pressure on the nerve in the carpal tunnel can cause numbness, tingling, burning or pain in the areas supplied by the nerve (part of the palm, the thumb, index, middle and ring finger) or weakness in the hand muscles. One or both hands may be affected.

Carpal Tunnel Syndrome is diagnosed through a thorough clinical examination. An accurate diagnosis is required to ensure the most appropriate treatment is implemented.

What symptoms might I experience?

- Numbness or a burning sensation in part of the palm, the thumb, index, middle and ring fingers.
- Waking at night with hand pain or numbness.
- Morning stiffness or cramping of the hand/wrist.
- Difficulty gripping objects.
- Difficulty using your hands (for example writing or turning a door knob).
- Thumb weakness.
- Mottled appearance of the hands.

What can I do if I have carpal tunnel syndrome?


- Avoid firmly gripping objects.

- Avoid repeated heavy lifting.
- For objects such as bags or clothes try to use your forearm instead of your hand when lifting - for example, rest the object on your arm rather than holding it in your hand.
- Avoid prolonged repetitive activities (like typing or stirring while cooking) or activities that aggravate your symptoms.
- Avoid sleeping with your arm and wrist tucked underneath you.

Things to try

- Ice your wrists for 15-20 minutes as required.
- Your physiotherapist may prescribe a wrist splint to hold the wrist in a neutral position, at night and/or during the day.
- Your physiotherapist may prescribe tendon gliding exercises. (See diagram)
 1. Keep the wrist straight, then bend the fingers like a hook (Step 1-2)
 2. Keep the wrist straight, then make a fist (Step 3-4)
 3. Keep the wrist straight, then bend the fingers keeping all finger joints straight. (Step 5-6) Repeat this with the fingers bent (Step 7-8) .

Note: repeat these exercises five times, three to four times per day.



Please only complete these exercises if your physiotherapist has instructed you to do so.

Will carpal tunnel syndrome go away after I've delivered?

In most cases symptoms will ease following delivery as pregnancy-related fluid retention reduces. Occasionally the symptoms are ongoing and can be aggravated by caring for your newborn baby. If your symptoms do not settle within the first few months after giving birth, please contact the Physiotherapy Department or your doctor for assistance with referral to a hand therapist.

Further Information

If you have any questions regarding this information, please contact Physiotherapy Department:

Mercy Hospital for Women Phone: 03 8458 4141

Werribee Mercy Hospital Phone: 03 8754 3150

Acknowledgements

Produced by: Physiotherapy Department

Date produced: July 2014

Date of last review: September 2019

Date for review: September 2021